



2018 CMS Web Interface

PREV-9 (NQF 0421): Preventive Care and Screening:
Body Mass Index (BMI) Screening and Follow-Up Plan

Measure Steward: CMS

Contents

INTRODUCTION 3

CMS WEB INTERFACE SAMPLING INFORMATION 4

BENEFICIARY SAMPLING 4

NARRATIVE MEASURE SPECIFICATION 5

DESCRIPTION: 5

IMPROVEMENT NOTATION: 5

INITIAL POPULATION: 5

DENOMINATOR: 5

DENOMINATOR EXCLUSIONS: 5

DENOMINATOR EXCEPTIONS: 5

NUMERATOR: 5

NUMERATOR EXCLUSIONS: 5

DEFINITIONS: 6

GUIDANCE: 6

SUBMISSION GUIDANCE 8

PATIENT CONFIRMATION 8

SUBMISSION GUIDANCE 9

DENOMINATOR CONFIRMATION 9

SUBMISSION GUIDANCE 10

NUMERATOR REPORTING 10

SUBMISSION GUIDANCE 11

NUMERATOR REPORTING 11

DOCUMENTATION REQUIREMENTS 11

APPENDIX I: PERFORMANCE CALCULATION FLOW: 12

APPENDIX II: DOWNLOADABLE RESOURCE MAPPING TABLE 19

APPENDIX III: MEASURE RATIONALE AND CLINICAL RECOMMENDATION STATEMENTS 20

RATIONALE: 20

CLINICAL RECOMMENDATION STATEMENTS: 21

APPENDIX IV: USE NOTICES, COPYRIGHTS, AND DISCLAIMERS 23

COPYRIGHT 23

INTRODUCTION

There are a total of 15 individual measures (including one composite consisting of two measures) included in the 2018 CMS Web Interface targeting high-cost chronic conditions, preventive care, and patient safety. The measures documents are represented individually and contain measure specific information. The corresponding coding documents are posted separately in an Excel format.

The measure documents are being provided to allow group practices and Accountable Care Organizations (ACOs) an opportunity to better understand each of the 15 individual measures included in the 2018 CMS Web Interface data submission method. Each measure document contains information necessary to submit data through the CMS Web Interface.

Narrative specifications, supporting submission documentation, and calculation flows are provided within each document. Please review all of the measure documentation in its entirety to ensure complete understanding of these measures.

CMS WEB INTERFACE SAMPLING INFORMATION

BENEFICIARY SAMPLING

For more information on the sampling process and methodology please refer to the 2018 CMS Web Interface Sampling Document, which will be made available during the performance year at CMS.gov.

NARRATIVE MEASURE SPECIFICATION

DESCRIPTION:

Percentage of patients aged 18 years and older with a BMI documented during the current encounter or during the previous twelve months AND with a BMI outside of normal parameters, a follow-up plan is documented during the encounter or during the previous twelve months of the current encounter

Normal Parameters: Age 18 years and older BMI => 18.5 and < 25 kg/m²

IMPROVEMENT NOTATION:

Higher score indicates better quality

INITIAL POPULATION:

All patients 18 and older on the date of the encounter with at least one eligible encounter during the measurement period

DENOMINATOR:

Equals Initial Population

DENOMINATOR EXCLUSIONS:

- Patients who are pregnant
- Patients who refuse measurement of height and/or weight or refuse follow-up

DENOMINATOR EXCEPTIONS:

Patients with a documented Medical Reason:

- Elderly Patients (65 or older) for whom weight reduction/weight gain would complicate other underlying health conditions such as the following examples:
 - Illness or physical disability
 - Mental illness, dementia, confusion
 - Nutritional deficiency, such as vitamin/mineral deficiency
- Patients in an urgent or emergent medical situation where time is of the essence and to delay treatment would jeopardize the patient's health status

NUMERATOR:

Patients with a documented BMI during the encounter or during the previous twelve months, AND when the BMI is outside of normal parameters, a follow-up plan is documented during the encounter or during the previous twelve months of the current encounter.

NUMERATOR EXCLUSIONS:

Not Applicable

DEFINITION:

BMI – Body mass index (BMI) is a number calculated using the Quetelet index: weight divided by height squared (W/H²) and is commonly used to classify weight categories. BMI can be calculated using:

- Metric Units: $BMI = \text{Weight (kg)} / (\text{Height (m)} \times \text{Height (m)})$

OR

- English Units: $BMI = \text{Weight (lbs.)} / (\text{Height (in)} \times \text{Height (in)}) \times 703$

Follow-Up Plan – Proposed outline of treatment to be conducted as a result of a BMI outside of normal parameters. A follow-up plan may include, but is not limited to: documentation of education, referral (for example a registered dietician, nutritionist, occupational therapist, physical therapist, primary care provider, exercise physiologist, mental health professional, or surgeon), pharmacological interventions, dietary supplements, exercise counseling or nutrition counseling.

GUIDANCE:

- There is no diagnosis associated with this measure.
- This measure is to be reported a minimum of once per performance period for patients seen during the performance period.
- This measure may be reported by eligible professionals who perform the quality actions described in the measure based on the services provided at the time of the qualifying visit and the measure-specific denominator coding.

BMI Measurement Guidance:

- Height and Weight - An eligible professional or their staff is required to measure both height and weight. Both height and weight must be measured within twelve months of the current encounter and may be obtained from separate encounters. Self-reported values cannot be used.
- The BMI may be documented in the medical record of the provider or in outside medical records obtained by the provider.
- If the most recent documented BMI is outside of normal parameters, then a follow-up plan is documented during the encounter or during the previous twelve months of the current encounter.
- If more than one BMI is reported during the measurement period, the most recent BMI will be used to determine if the performance has been met.
- Review the exclusions criteria to determine those patients that BMI measurement may not be appropriate or necessary.

Follow-Up Plan Guidance:

- The documented follow-up plan must be based on the most recent documented BMI, outside of normal parameters, example: "Patient referred to nutrition counseling for BMI above or below normal parameters." (See Definitions for examples of follow-up plan treatments).

Variation has been noted in studies exploring optimal BMI ranges for the elderly (see Donini et al., (2012); Holme and Tonstad (2015); and Diehr et al. (2008). Notably however, all these studies have arrived at ranges that differ from the standard range for ages 18 and older, which is ≥ 18.5 and < 25 kg/m². For instance, both Donini et al. (2012) and Holme and Tonstad (2015) reported findings that suggest that higher BMI (higher than the upper end of 25kg/m²) in the elderly may be beneficial. Similarly, worse outcomes have been associated with being underweight (at a threshold higher than 18.5 kg/m²) at age 65 (Diehr et al. 2008). Because of optimal BMI range variation recommendations from these studies, no specific optimal BMI range for the elderly is used. However, it may be

appropriate to exempt certain patients from a follow-up plan by applying the exception criteria. Review the following to apply the Medical Reason exception criteria:

The Medical Reason exception could include, but is not limited to, the following patients as deemed appropriate by the health care provider:

- Elderly Patients (65 or older) for whom weight reduction/weight gain would complicate other underlying health conditions such as the following examples:
 - Illness or physical disability
 - Mental illness, dementia, confusion
 - Nutritional deficiency such as vitamin/mineral deficiency
- Patients in an urgent or emergent medical situation where time is of the essence and to delay treatment would jeopardize the patient's health status

SUBMISSION GUIDANCE

PATIENT CONFIRMATION

Establishing patient eligibility for submitting requires the following:

- Determine if the patient's medical record can be found
 - If you can locate the medical record select "Yes"

OR

- If you cannot locate the medical record select "No - Medical Record Not Found"

OR

- Determine if the patient is qualified for the sample
 - If the patient is deceased, in hospice, moved out of the country or was enrolled in HMO select "Not Qualified for Sample", select the applicable reason from the provided drop-down menu, and enter the date the patient became ineligible

Guidance Patient Confirmation

If "No – Medical Record Not Found" or "Not Qualified for Sample" is selected, the patient is completed but not confirmed. The patient will be "skipped" and another patient must be submitted in their place, if available. The CMS Web Interface will automatically skip any patient for whom "No – Medical Record Not Found" or "Not Qualified for Sample" is selected in all other measures into which they have been sampled.

If "Not Qualified for Sample" is selected and the date is unknown, you may enter the last date of the measurement period (i.e., 12/31/2018).

The Measurement Period is defined as January 1 – December 31, 2018.

NOTE:

- ***In Hospice:** Select this option if the patient is not qualified for sample due to being in hospice care at any time during the measurement period (this includes non-hospice patients receiving palliative goals or comfort care)*
- ***Moved out of Country:** Select this option if the patient is not qualified for sample because they moved out of the country any time during the measurement period*
- ***Deceased:** Select this option if the patient died during the measurement period*
- ***HMO Enrollment:** Select this option if the patient was enrolled in an HMO at any time during the measurement period (i.e., Medicare Advantage, non-Medicare HMOs, etc.)*

SUBMISSION GUIDANCE

DENOMINATOR CONFIRMATION

- Determine if the patient is qualified for the measure
 - If the patient is qualified for the measure select "Yes"

OR

- If there is a denominator exclusion for patient disqualification from the measure select "[Denominator Exclusion](#)"

OR

- If there is an "other" CMS approved reason for patient disqualification from the measure select "No-Other CMS Approved Reason"

Denominator Exclusion codes can be found in the 2018 CMS Web Interface PREV Coding Document. The Downloadable Resource Mapping Table can be located in Appendix II of this document.

Guidance *Denominator*

If "Denominator Exclusion" or "No – Other CMS Approved Reason" is selected, the patient will be "skipped" and another patient must be submitted in their place, if available. The patient will only be removed from the measure for which one of these options was selected, not all CMS Web Interface measures.

CMS Approved Reason may only be selected when approved by CMS. To request a CMS Approved Reason, you would need to provide the patient rank, measure, and reason for request in a Quality Payment Program Service Desk inquiry. A CMS decision will be provided in the resolution of the inquiry. Patients for whom a CMS Approved Reason is selected will be "skipped" and another patient must be submitted in their place, if available.

NOTE:

- *The timing components for the Denominator Exclusions are as follows:*
 - *Patients who are pregnant any time during the measurement period*
 - *Patient refusal could occur at any encounter [during the measurement period](#)*

SUBMISSION GUIDANCE

NUMERATOR SUBMISSION

- Determine if the patient had a BMI documented during the most recent visit or in the last 12 months prior to the most recent visit
- If the patient has not had a BMI documented select "No"

OR

- If the patient has had a BMI calculated select "Yes"

Numerator codes can be found in the 2018 CMS Web Interface PREV Coding Document. The Downloadable Resource Mapping Table can be located in Appendix II of this document.

Guidance Numerator

NOTE:

- *The twelve month look-back can be calculated as a look-back of less than 13 months from the most recent encounter*
- *The twelve month look-back could potentially fall outside the measurement year*

SUBMISSION GUIDANCE

NUMERATOR SUBMISSION

- If the patient had a BMI calculated, determine if the most recent BMI is within normal parameters
 - If the most recent BMI is outside of normal parameters select "No"

IF NO

- If the patient's most recent BMI was not within normal limits, determine if a follow-up plan was documented
 - If there was no follow-up plan documented select "No"

OR

- If there was a follow-up plan documented select "Yes"

OR

- When a recommended follow-up for an abnormal BMI is not documented for medical reasons select "No - [Denominator Exception](#) - Medical Reasons"

OR

- If the most recent BMI is within normal parameters select "Yes"

Numerator and Denominator Exception codes can be found in the 2018 CMS Web Interface PREV Coding Document. The Downloadable Resource Mapping Table can be located in Appendix II of this document.

Guidance Numerator

NOTE:

- *Amputees are not considered denominator exceptions*
- *BMI calculation and recommended follow-up plan cannot be completed during a telehealth encounter*
- *The timing component for the Denominator Exception is the date of the encounter with the calculated BMI or within the previous 12 months of the current encounter*


DOCUMENTATION REQUIREMENTS

When submitting data through the CMS Web Interface, the expectation is that medical record documentation is available that supports the action submitted in the CMS Web Interface i.e., medical record documentation is necessary to support the information that has been submitted.

Appendix I: Performance Calculation Flow:

Patient Confirmation Flow


For 2018, confirmation of the "Medical Record Found", or indicating the patient is "Not Qualified for Sample" with a reason of "In Hospice", "Moved out of Country", "Deceased", or "HMO Enrollment", will only need to be done **once** per patient.



*See the Measure Submission Document for further instructions on how to submit this measure
 **If date is unknown, enter 12/31/2018

Measure Confirmation Flow for PREV-9

For 2018, measure specific reasons a patient is "Not Confirmed" or excluded for "Denominator Exclusion" or "Other CMS Approved Reason" will need to be done for each measure where the patient appears.




*See the Measure Submission Document for further instructions on how to submit this measure

**Further information regarding patient selection for specific disease and patient care measures can be found in the CMS Web Interface Sampling Methodology Document. For patients who have the incorrect date of birth listed, a change of the patient date of birth by the abstractor may result in the patient no longer qualifying for the PREV-9 measure. If this is the case, the system will automatically remove the patient from the measure requirements.

***"Other CMS Approved Reason" may only be selected if you have received an approval from CMS in the resolution of a requested Quality Payment Program Service Desk Inquiry at qpp@cms.hhs.gov

Measure Flow for PREV-9



The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used as a substitution for the measure specification. For Downloadable Resource Mapping Table, go to Appendix II and use the Variable Names located in the appendix along with the applicable tabs within the PREV Code Document.

SAMPLE CALCULATION:

Performance Rate=

$$\frac{\text{Performance Met (a}^1\text{=85 Patients + a}^2\text{=90 Patients)}}{\text{Denominator (d=222 Patients) - Denominator Exception (b=25 Patients)}} = \frac{175 \text{ Patients}}{197 \text{ Patients}} = 88.83\%$$

CALCULATION MAY CHANGE PENDING PERFORMANCE MET ABOVE

*See the Measure Submitting Document for further instructions on how to submit this measure

Patient Confirmation Flow

For 2018, confirmation of the "Medical Record Found", or indicating the patient is "Not Qualified for Sample" with a reason of "In Hospice", "Moved out of Country", "Deceased", or "HMO Enrollment", will only need to be done **once** per patient. Refer to the Measure Submission Document for further instructions.

1. Start Patient Confirmation Flow.
2. Check to determine if Medical Record can be found.
 - a. If no, Medical Record not found, mark appropriately for completion and stop abstraction. This removes the patient from the beneficiary sample for all measures. The patient will be skipped and replaced. Stop processing.
 - b. If yes, Medical Record found, continue processing.
3. Check to determine if Patient Qualified for the sample.
 - a. If no, the patient does not qualify for the sample, select the reason why and enter the date (if date is unknown, enter 12/31/2018) the patient became ineligible for sample. For example; In Hospice, Moved out of Country, Deceased, HMO Enrollment. Mark appropriately for completion and stop abstraction. This removes the patient from the beneficiary sample for all measures. The patient will be skipped and replaced. Stop processing.
 - b. If yes, the patient does qualify for the sample; continue to the Measure Confirmation Flow for PREV-9.

Measure Confirmation Flow for PREV-9

For 2018, measure specific reasons a patient is "Not Confirmed" or excluded for "Denominator Exclusion" or "Other CMS Approved Reason" will need to be done for each measure where the patient appears. Refer to the Measure Submission Document for further instructions.

1. Start Measure Confirmation Flow for PREV-9. Complete for consecutively ranked patients aged 18 years and older at the beginning of the measurement period. Further information regarding patient selection for specific disease and patient care measures can be found in the CMS Web Interface Sampling Methodology Document. For patients who have the incorrect date of birth listed, a change of the patient date of birth by the abstractor may result in the patient no longer qualifying for the PREV-9 measure. If this is the case, the system will automatically remove the patient from the measure requirements.
2. Check to determine if the patient qualifies for the measure (Denominator Exclusion).
 - a. If no, the patient does not qualify for the measure select: Denominator Exclusion for patient disqualification. Mark appropriately for completion and stop abstraction. Patient is removed from the performance calculations for this measure. The patient will be skipped and replaced. Stop processing.
 - b. If yes, the patient does qualify for the measure, continue processing.
3. Check to determine if the patient qualifies for the measure (Other CMS Approved Reason).
 - a. If no, the patient does not qualify for the measure select: No – Other CMS Approved Reason for patient disqualification. Mark appropriately for completion and stop abstraction. Patient is removed from the performance calculations for this measure. The patient will be skipped and replaced. "Other CMS Approved Reason" may only be selected if you have received an approval from CMS in the resolution of a requested Quality Payment Program Service Desk Inquiry at qpp@cms.hhs.gov. Stop Processing.
 - b. If yes, the patient does qualify for the measure, continue to the PREV-9 measure flow.

Measure Flow for PREV-9

The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used as a substitution for the measure specifications. For Downloadable Resource Mapping Table, go to Appendix II and use the Variable Names located in the appendix along with the applicable tabs within the PREV Coding Document.

1. Start processing 2018 PREV-9 (NQF 0421) Flow for the patients that qualified for sample in the Patient Confirmation Flow and the Measure Confirmation Flow for PREV-9. Note: Include remainder of patients listed in CMS Web Interface that were consecutively confirmed and completed for this measure in the denominator. For the sample calculation in the flow these patients would fall into the 'd' category (eligible denominator, i.e. 222 patients).
2. Check to determine if the patient had a BMI calculated during the most recent visit or within the last twelve months prior to the most recent visit.
 - a. If no, the patient did not have a BMI calculated during the most recent visit or within the last twelve months prior to the most recent visit, performance is not met and should not be included in the numerator. Stop processing.
 - b. If yes, the patient had a BMI calculated during the most recent visit or within the last twelve months prior to the most recent visit, continue processing.
3. Check to determine if the patient's most recent BMI is within normal parameters.
 - a. If no, the patient's most recent BMI is not within normal parameters, continue processing.
 - b. If yes, the patient's most recent BMI is within normal parameters, performance is met and the patient will be included in the numerator. For the sample calculation in the flow these patients would fall into the 'a¹' category (numerator, i.e. 85 patients). Stop processing.
4. Check to determine if the patient's follow-up plan was documented.
 - a. If no, the patient's follow-up plan was not documented, continue processing.
 - b. If yes, the patient's follow-up plan was documented, performance is met and the patient will be included in the numerator. For the sample calculation in the flow these patients would fall into the 'a²' category (numerator, i.e. 90 patients). Stop processing.
5. Check to determine if the patient's follow-up plan was Not documented for a denominator exception, medical reason(s).
 - a. If no, the patient's follow-up plan was Not documented for a denominator exception, medical reason(s), performance is not met and the patient should not be included in the numerator. Stop processing.
 - b. If yes, the patient's follow-up plan was Not documented for a denominator exception, medical reason(s), this is a denominator exception and the case should be subtracted from the denominator. For the sample calculation in the flow these patients would fall into the 'b' category (denominator exception, i.e. 25 patients). Stop processing.

Sample Calculation

Performance Rate Equals

Performance Met is category 'a¹ plus a²' in the measure flow (175 patients)

Denominator is category 'd' in the measure flow (222 patients)

Denominator Exception is category 'b' in the measure flow (25 patients)

175 (Performance Met) divided by 197 (Denominator minus Denominator Exception) equals a performance rate of
88.83 percent

Calculation May Change Pending Performance Met

Appendix II: Downloadable Resource Mapping Table

Each data element within this measure's denominator or numerator is defined as a pre-determined set of clinical codes. These codes can be found in the 2018 CMS Web Interface PREV Coding Document.

*PREV-9: Preventive Care and Screening: Screening for Body Mass Index (BMI) Screening and			
Measure Component/Excel Tab	Data Element	Variable Name	Coding System(s)
Denominator Exclusion/ Denominator Exclusion Codes	Exclusion	PREGNANCY_CODE	I10 SNM
		PATIENT_REASON_REFUSED	SNM
Numerator/Numerator Codes/Numerator Drug Codes	BMI Documented	BMI_CODE	LN
		BMI_ABNORMAL_CODE	HCPCS
	BMI Abnormal	BMI_ABNORMAL_CODE	HCPCS
		OVERWEIGHT_CODE	SNM
		UNDERWEIGHT_CODE	SNM
	Follow-up Plan	BMI_ABNORMAL_CODE	HCPCS
		BMI_FOLLOW_UP_CODE	I10 C4 HCPCS SNM
		REFERRAL_CODE	SNM
		BMI_DRUG_CODE	RxNorm (Drug EX=N)
	Denominator Exception/ Denominator Exception Codes	Medical Reason	MEDICAL_OTHER_REASON

** For EHR mapping, the coding within PREV-9 is considered to be all inclusive*

Appendix III: Measure Rationale and Clinical Recommendation Statements

RATIONALE:

BMI Above Normal Parameters

Obesity is a chronic, multifactorial disease with complex psychological, environmental (social and cultural), genetic, physiologic, metabolic and behavioral causes and consequences. The prevalence of overweight and obese people is increasing worldwide at an alarming rate in both developing and developed countries. Environmental and behavioral changes brought about by economic development, modernization and urbanization have been linked to the rise in global obesity. The health consequences are becoming apparent (ICSI 2013. p.6).

Nationally, nearly 38 percent of adults are obese [NHANES, 2013-2014 data]. Nearly 8 percent of adults are extremely obese (BMI greater than or equal to 40.0); Obesity rates are higher among women (40.4 percent) compared to men (35.0 percent). Between 2005 and 2014, the difference in obesity among women was 5.1 percent higher among women and 1.7 percent higher among men. Women are also almost twice as likely (9.9 percent) to be extremely obese compared to men (5.5 percent); In addition, rates are the highest among middle-age adults (41 percent for 40- to 59-year-olds), compared to 34.3 percent of 20- to 39-year-olds and 38.5 percent of adults ages 60 and older (Flegal KM, Kruszon-Moran D, Carroll MD, et al, 2016, p.2286-2290)..

Obesity is one of the biggest drivers of preventable chronic diseases and healthcare costs in the United States. Currently, estimates for these costs range from \$147 billion to nearly \$210 billion per year (Cawley J and Meyerhoefer C., 2012 & Finkelstein, Trogon, Cohen, et al., 2009). There are significant racial and ethnic inequities [NHANES, 2013-2014 data]: Obesity rates are higher among Blacks (48.4 percent) and Latinos (42.6 percent) than among Whites (36.4 percent) and Asian Americans (12.6 percent). The inequities are highest among women: Blacks have a rate of 57.2 percent, Latinos of 46.9 percent, Whites of 38.2 percent and Asians of 12.4 percent. For men, Latinos have a rate of 37.9 percent, Blacks of 38.0 percent and Whites of 34.7 percent. Black women (16.8 percent) are twice as likely to be extremely obese as White women (9.7 percent) (Flegal KM, Kruszon-Moran D, Carroll MD, et al., 2016, pp. 2284-2291).

BMI continues to be a common and reasonably reliable measurement to identify overweight and obese adults who may be at an increased risk for future morbidity. Although good quality evidence supports obtaining a BMI, it is important to recognize it is not a perfect measurement. BMI is not a direct measure of adiposity and as a consequence it can over- or underestimate adiposity. BMI is a derived value that correlates well with total body fat and markers of secondary complications, e.g., hypertension and dyslipidemia (Barlow, 2007).

In contrast with waist circumference, BMI and its associated disease and mortality risk appear to vary among ethnic subgroups. Female African American populations appear to have the lowest mortality risk at a BMI of 26.2-28.5 kg/m² and 27.1-30.2 kg/m² for women and men, respectively. In contrast, Asian populations may experience lowest mortality rates starting at a BMI of 23 to 24 kg/m². The correlation between BMI and diabetes risk also varies by ethnicity (LeBlanc, 2011. p.2-3).

Screening for BMI and follow-up therefore is critical to closing this gap and contributes to quality goals of population health and cost reduction. However, due to concerns for other underlying conditions (such as bone health) or nutrition related deficiencies providers are cautioned to use clinical judgment and take these into account when considering weight management programs for overweight patients, especially the elderly (NHLBI Obesity Education Initiative, 1998, p. 91).

BMI below Normal Parameters

On the other end of the body weight spectrum is underweight (BMI <18.5 kg/m²), which is equally detrimental to population health. When compared to normal weight individuals (BMI 18.5-25 kg/m²), underweight individuals have

significantly higher death rates with a Hazard Ratio of 2.27 and 95% confidence intervals (CI) = 1.78, 2.90 (Borrell & Lalitha (2014).

Poor nutrition or underlying health conditions can result in underweight (Fryer & Ogden, 2012). The National Health and Nutrition Examination Survey (NHANES) results from the 2007-2010 indicate that women are more likely to be underweight than men (2012). Therefore patients should be equally screened for underweight and followed up with nutritional counselling to reduce mortality and morbidity associated with underweight.

CLINICAL RECOMMENDATION STATEMENTS:

As cited in Fetch et al. (2013), The Institute for Clinical Systems Improvement (ICSI) Health Care Guideline, Prevention and Management of Obesity for Adults provides the Strength of Recommendation as Strong for the following:

- Record height, weight and calculate body mass index at least annually
- Clinicians should consider waist circumference measurement to estimate disease risk for patients who have normal or overweight BMI scores. For adult patients with a BMI of 25 to 34.9 kg/m², sex-specific waist circumference cutoffs should be used in conjunction with BMI to identify increased disease risk.

Individuals who are overweight (BMI 25<30), and who do not have indicators of increased CVD risk (e.g., diabetes, pre-diabetes, hypertension, dyslipidemia, elevated waist circumference) or other obesity-related comorbidities and individuals who have a history of overweight and are now normal weight with risk factors at acceptable levels:

"Advise to frequently measure their own weight, and to avoid weight gain by adjusting their food intake if they start to gain more than a few pounds. Also, advise patients that engaging in regular physical activity will help them avoid weight gain." (2013 AHA/AAC/TOS Obesity Guideline, p. S113)

"Advise overweight and obese individuals who would benefit from weight loss to participate for ≥6 months in a comprehensive lifestyle program that assists participants in adhering to a lower calorie diet and in increasing physical activity through the use of behavioral strategies... NHLBI Grade A (Strong)" (2013 AHA/AAC/TOS Obesity Guideline, p. S109)

USPSTF Clinical Guideline (Grade B Recommendation)

Individuals with a body mass index (BMI) of 30 kg/m² or higher should be offered or referred to intensive, multicomponent behavioral interventions that include the following components:

- Behavioral management activities, such as setting weight-loss goals
- Improving diet or nutrition and increasing physical activity
- Addressing barriers to change
- Self-monitoring
- Strategizing how to maintain lifestyle changes

Nutritional safety for the elderly should be considered when recommending weight reduction. "A clinical decision to forego obesity treatment in older adults should be guided by an evaluation of the potential benefits of weight reduction for day-to-day functioning and reduction of the risk of future cardiovascular events, as well as the patient's motivation for weight reduction. Care must be taken to ensure that any weight reduction program minimizes the likelihood of adverse effects on bone health or other aspects of nutritional status" Evidence Category D. (NHLBI Obesity Education Initiative, 1998, p. 91). In addition, weight reduction prescriptions in older persons should be accompanied by proper nutritional counseling and regular body weight monitoring. (NHLBI Obesity Education Initiative, 1998, p. 91).

The possibility that a standard approach to weight loss will work differently in diverse patient populations must be considered when setting expectations about treatment outcomes. Evidence Category B. (NHLBI Obesity Education Initiative, 1998).

Appendix IV: Use Notices, Copyrights, and Disclaimers

COPYRIGHT

Limited proprietary coding is contained in the measure specifications for convenience. Users of the proprietary code sets should obtain all necessary licenses from the owners of these code sets. Quality Insights, Inc. disclaims all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specifications.

(CPT®) contained in the Measure specifications is copyright 2007-2017 American Medical Association.

LOINC® copyright 2004-2017 [2.58] Regenstrief Institute, Inc. This material contains SNOMED Clinical Terms (SNOMED CT®) copyright 2004-2017 [2017-03] International Health Terminology Standards Development Organization. All Rights Reserved.

THE MEASURES AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND.